

Intermediate ECG Course – Part 2

March-April 2010

Joe M. Moody, Jr, MD


UTHSCSA and STVAHCS

Topics in Intermediate ECG

- Consolidation of prior information with additional details
- Not “advanced”, but feel free to ask advanced questions
- Causes of axis deviation and wide QRS (1)
- Infarction and causes of ST segment shifts (2)
- Electrolyte effects on the ECG (3)
- Flutter versus fib, and ventricular patterns (4)
- AV conduction and AV dissociation (4)
- Tachyarrhythmias, wide and narrow QRS (5, 6)
- Integrating ECG and clinical information (7,8)


Ischemia, Injury, Infarction


- Context = resting ECG
- Ischemia: T wave inversion in distribution of affected leads (does localize)
- Injury:
 - Subendocardial injury: ST segment depression (does not localize)
 - Subepicardial/transmural injury: ST segment elevation in affected leads (does localize)
- Infarction: Q wave formation in affected leads (does localize)



ECG-Ischemia

T wave inversion







ST depression: subendocardial injury pattern


ECG Injury

ST elevation: transmural (or subepicardial) injury pattern




Ischemia, Injury, Infarction - 3

- Infarction - irreversible loss of myocardium, replaced eventually by electrically silent collagen
 - ECG - initial forces directed away from area of infarction, with abnormal Q wave in affected leads
 - Abnormal Q (Pathologic Q) - 0.03 sec wide and depth of 1 mm in 2 contiguous leads (not III)



Ischemia, Injury, Infarction - 3

- Infarction - irreversible loss of myocardium, replaced eventually by electrically silent collagen
 - ECG - initial forces directed away from area of infarction, with abnormal Q wave in affected leads
 - Abnormal Q (Pathologic Q) - 0.03 sec wide and depth of 1 mm in 2 contiguous leads (not III)



Infarction Location

Posterior!!!

Lateral,
High Lateral

aVR

Right Precordial,
Septal

V4

II

aVL

V2

V5

III
Inferior

aVF

V3

Antero-
Lateral

V6

Lateral – diagonal branch of LAD
Inferior – RCA or LCX
Septal – LAD or septal perforator
Anterior – Mid LAD
Anterolateral – LAD with diagonal


Mid Precordial,
Anterior,
Localized Anterior

Causes of Prominent Anterior Forces ($R > S$ in V1)

- Normal variant in young adult
- RVH
- True posterior infarction
- Pulmonary disease with displacement of the heart
- WPW pattern (“type A”)
- Duchenne’s muscular dystrophy
- Lead misplacement
- Dextrocardia


Causes of ST Segment Elevation

- Acute transmyocardial injury
- Hyperkalemia
- Pericarditis
- Normal variant, early repolarization
- Hypothermia (Osborn wave)
- Brugada Syndrome
- Less common
 - Pulmonary embolism and acute cor pulmonale
 - Cardiac tumor
 - Aortic dissection
 - Post mitral valvuloplasty
 - Pancreatitis and GB disease
 - Myocarditis
 - Septic shock
 - Anaphylactic reaction
 - Drug overdose (tricyclic, etc)



ECG - Pericarditis

- = Diffuse ST elevation
- = PR segment depression
- = No reciprocal changes except aVR



Time Course of Infarction

- Time course is highly affected by acute intervention (“interruption of the infarction”)
- Hyperacute T waves – minutes, generally less than 10 minutes, usually missed
- ST elevation – onset in seconds to minutes, decreases markedly during the first 7-12 hours, usually gone in few days, represents aneurysm if present more than 8 weeks
- Q wave appears within 6-12 hours (9) and is usually permanent
- T wave inversion onset is after ST elevation and may persist several days, weeks, months, or years

Arrhythmia in Inferior Wall MI

- Usually bradycardia, sinus bradycardia (SA node may be affected, also vagal effect)
- Possible ventricular tachyarrhythmias
- First degree AV block or second degree Mobitz I (Wenckebach) AV block or third degree AV block with narrow QRS escape rhythm (AV Node branch is from the RCA or the LCX); AV block is usually transient and resolves spontaneously

Arrhythmia in Anterior Wall MI

- Sinus tachycardia (MI is large resulting in poor stroke volume so increased HR to maintain cardiac output)
- Ventricular tachyarrhythmias
- Conduction abnormalities are usually below the AV node in the bundle branches
- Second degree AV block may be Mobitz II
- AV block usually has a wide escape rhythm and is usually permanent


Conduction Problems

- AVN delay in inferior MI and proximal RCA, frequently with RV involvement
- RBBB +/- AFB indicates proximal LAD
- AFB in Inferior MI indicates LAD disease

Clinical Presentation of MI in Different Vessel Involvement


- LAD: large MIs, anterior lateral septal inferoapical and proximal bundle branches
- RCA: sinus node in 55%, RV, AVN, posteromedial papillary muscle, inferior LV and variably posterior and lateral; vagal activation, sinus bradycardia, RV involvement, papillary muscle dysfunction and MI
- LCX: posterior and variably inferior and lateral ... abnormalities in second half of the QRS complex and frequently cause underestimation of the area at risk and undertreatment of the patient

September 5, 2004 21:33




Woman less than 50 yo.

September 6, 2004 05:36



Woman less than 50 yo.

Anterior MI



Localization of Anterior MI Lesion by ECG

- 100 pts, first anterior MI, tracing with most pronounced ST change before reperfusion therapy; ST elev >2mm in V2 and V3
- Excluded LBBB, prior CABG, LVH, prior cardiac surgery
- ST elev in aVR in 43% prox to S1 and 5% dist to S1, 0.4 (0.2-1.8mm)

ECG in Anterior MI - ST elevation in V2-4

- 40% are proximal to 1SP and D1
 - ST elevation in aVR and ST elev >2.5 mm in V1, ST depression in 2, 3, and F and maybe V5-6, and Q in aVL
- 40% are distal to both
 - Dominance of inferoapical area, absent ST depression and maybe elevation in 2, 3, and F, sometimes Qs in V4-6,
- 10% are proximal to D1 but not 1SP
 - Q in left lateral leads, ST depression in III, and no ST depression in II
- 10% are proximal to 1SP but not D1
 - ST elevation in aVR and >2.5 mm elevation in V1, and ST depression in V5, and maybe ST elevation in V3R, and ST depression in aVL, a very specific finding, also ST elevation in 2, 3, and F

Site of Occlusion of LAD

Criterion	Occlusion Site	Sens	Spec	PPA	NPA
CRBBB	Prox to S1	12	100	100	62
ST \uparrow V1>2.5	Prox S1	12	100	100	61
ST \uparrow aVR	Prox S1	43	95	86	70
ST \downarrow V5	Prox S1	17	98	88	62
Q in L	Prox D1	44	85	67	69
ST \downarrow II>1.0 mm	Prox S1/D1	34	98	93	68
Q V5	Dist S1	24	93	71	53
ST \downarrow in L	Dist D1	22	95	87	46
No ST \downarrow III	Dist S1/D1	41	95	92	53


Localizing LAD Lesion

Table 2. Electrocardiographic Predictors of Left Anterior Descending Coronary Artery (LAD) Occlusion Proximal to the First Septal Perforator (S1) and/or the First Diagonal Branch (D1)

Predictors of LAD Occlusion Proximal to S1						
	Sens	Spec	PPV	NPV	LR	p-Value
ST \uparrow _{aVR}	43	95	86	70	8.6	0.000
ST \downarrow _{II} ≥ 1.0 mm	36	100	100	68		0.000
ST \downarrow _{III} ≥ 1.0 mm	60	71	60	71	2.1	0.002
ST \downarrow _{III} ≥ 2.5 mm	33	97	88	67	11.0	0.000
ST \downarrow _{aVF} ≥ 1.0 mm	52	84	71	71	3.3	0.000
ST \downarrow _{aVF} ≥ 2.0 mm	26	97	85	64	8.7	0.002
cRBBB	14	100	100	62		0.004
ST \downarrow _{V5}	17	98	88	62	8.5	0.009
ST \uparrow _{V1} > 2.5 mm	12	100	100	61		0.011
Predictors of LAD Occlusion Proximal to D1						
ST \downarrow _{II} ≥ 1.0 mm	34	98	93	68	17.0	0.000
ST \downarrow _{III} ≥ 1.0 mm	66	75	64	76	2.6	0.000
ST \downarrow _{III} ≥ 2.5 mm	32	95	81	67	6.4	0.001
ST \downarrow _{aVF} ≥ 1.0 mm	54	85	71	72	3.6	0.000
ST \downarrow _{aVF} ≥ 2.0 mm	27	97	85	66	9.0	0.001
Q _{aVL}	44	85	67	69	2.9	0.002

cRBBB = complete right bundle branch block; LR = likelihood-ratio; NPV = negative predictive value; PPV = positive predictive value; sens = sensitivity; spec = specificity; ST \downarrow = ST-depression; ST \uparrow = ST-elevation; Q = abnormal Q-wave.

Inferior Infarction



Inferior MI, Naming the Artery


- Simple inferior MI – RCA
- Inferior MI with RV involvement – RCA
- Inferior MI with posterior involvement – LCX
- Inferior MI with lateral involvement - LCX

RCA vs LCX


- RCA STel in III>II (STdep aVL>I)
- LCX STel in II>III (I with ST 0 or elev)
- STEL in V5-V6 of little value in differentiating, just larger area at risk
- V4R: Proximal RCA ST elev in V4R and upright T wave, distal RCA no ST elev and upright T wave, LCX no ST elev and inverted T wave (reliability requires significant ST elevation in inferior leads)
- Anterior ST depression: may extend V1-V6, if maximal in V4-6 likely 3-V dz and lower EF, absence indicates RCA and presence unhelpful to distinguish; maximal ST depression in V2-3 predicts LCX
- Isolated RVMI, minor changes inferiorly, ST elevation in V1-2 and V3-4R (I've also heard that if V1>V2, indicates RVMI)

RIGHT CORONARY ARTERY MI

CIRCUMFLEX CORONARY ARTERY MI



Inferoseptal



Posterolateral

**ST SEGMENT VECTOR IN RIGHT CORONARY ARTERY
VS CIRCUMFLEX MYOCARDIAL INFARCTION**

FIGURE 53-9 Schematic presentation of the ST-segment vector with inferoposterior MI caused by a right coronary artery (RCA) or circumflex coronary artery (CX). As shown, RCA occlusion leads to predominant ischemia in the inferoseptal area with an ST-segment vector pointing toward lead III. In CX occlusion, the ischemic area is located posterolaterally, resulting in an ST-segment vector directed toward lead II.




FIGURE 53-11 Diagram showing the coronary arteries and the possible sites of coronary artery occlusion leading to inferoposterior MI. In the right coronary artery (RCA), the occlusion may be before (proximal) the right ventricular (RV) branch or after it (distal). As shown in proximal RCA occlusion, the RV is involved in the MI.

VALUE OF ST-T SEGMENT CHANGES IN LEAD V_{4R} IN ACUTE INFERO-POSTERIOR MYOCARDIAL INFARCTION

**ST \uparrow \geq 1 mm
POS T-WAVE**



PROXIMAL OCCLUSION RCA

**NO ST \uparrow :
POS T-WAVE**



DISTAL OCCLUSION RCA

NEG T-WAVE



OCCLUSION CX

FIGURE 53-12 Characteristic ST-T-segment changes in lead V_{4R} in cases of proximal RCA, a distal RCA occlusion, or a CX occlusion (see text).

LCX Occlusion

- 84 patients: 2 normal, 35 with Q waves, 43 true posterior MI, 2 ST-T abnormalities, 2 LBBB
- Inferior Q: peripheral stenoses
- Lateral Q: central stenoses
- True posterior both central and peripheral

RV Myocardial Infarction

- Cause: proximal RCA lesion
- Hemodynamics: low BP, elevated JVP, clear lungs (RV can't get blood across to the LV), may have severe hypotension with NTG, RA pressure equal or greater than PAW
- ECG indicators: ST elevation in V1 greater than V2, or RV4 ST elevation of at least 1 mm

LMCA disease

- Severe hemodynamic deterioration
- Subtotal with collat from RCA is more common as UA with marked ST depression in I, II, and V4-V6 and ST elev in aVR
- Total LMCA obst aVR ST elevation >0.05 mV in 88% as opposed to LAD 43% or RCA 8%, and higher amplitude 0.16mV

Atrial infarction

- Atrial repolarization: elevation in I, II, III, V5 or V6, or depression in precordial leads
- May be seen in 10% inferoposterior MI
- Indicates proximal occlusion of RCA or LCX
- Complication atrial fibrillation or flutter, MAR, sinus arrest, atrial rupture (rare)

Caveats in ECG Diagnosis of MI

- There are many exceptions
- Most reliable with first MI
- Impaired with multivessel disease, prior MI, collateral circulation, LBBB, preexcitaion and paced rhythms

Pseudo-Infarction

- Loss of Viable Myocardium
- Altered distribution of myocardial mass
- Altered sequence of depolarization or abnormal cardiac position

Pseudo-Infarction

- Loss of Viable Myocardium:
 - Duchenne-type muscular dystrophy - inferoposterior MI pattern but less wide Q waves, also
 - Friedrich's ataxia
- Altered distribution of myocardial mass:
 - HCM,
 - LVH,
 - RVH


Pseudo-Infarction

- Altered sequence of depolarization or abnormal cardiac position: stocky people inferior MI, poor R progression in
 - R to L septal depolarization: incomplete LBBB, LVH, dextrocardia, “corrected” TGA
 - Inferior deviation of initial QRS force (LAFB, thin persons)
 - Downward displacement of origin of initial QRS (emphysema)
 - Position of electrodes (posterior pericardial effusion, high electrodes)
 - Pseudo-Q wave with perpendicular orientation of initial QRS deflection to the lead axis

More Pseudo-Infarction

- Pneumothorax, scoliosis
- Pulmonary embolism and acute cor pulmonale
- Hyperkalemia
- Q waves in PVCs

Practice 1



Rhythm sinus with irregularities


Dramatic ST elevation in inferior and lateral precordial leads

Significant ST depression in right precordial leads

Early (marked ST change and no T evolution, not much Q wave)

Acute inferoposterolateral MI

Practice 2



Sinus rhythm, normal PR interval

Insignificant inferior Q waves


Diffuse T wave abnormality

Significant anterior T wave inversion

In the setting of unstable angina, this finding is anterior ischemia

These “Wellens T waves” indicate LAD disease

Practice 3



Sinus rhythm, rate near 100


Anterior MI, old, with R wave decrement from V2 to V3

Possible anterolateral involvement, because R waves in V4-6 are small

Q in III is inconsequential if in isolation, but here has a Q in aVF

Association of localized anterior and inferior may indicate “apical”

Practice 4



Sinus rhythm rate about 80

Significant Qs in II, III, and aVF, with a little ST elevation and


With no T wave inversion, but also

With prolonged QT interval

I guess this inferior MI is probably acute or recent but

I'd want good clinical correlation

Practice 5



Sinus tachycardia rate 150...(be sure it is not atrial flutter)


Significant diffuse ST depression in 9 leads

Subendocardial ischemia

No infarction Q waves


Does not look like hypertrophy

Practice 6




Sinus rhythm, one PAC, very acute inferoposterolateral MI

Practice 6B




Same patient, now not necessarily sinus
RBBB pattern has developed – is this rate-related or VT?
I find no P waves, rate about 130

Practice 6C




Back to sinus, still very early, but ST elevation is less

Practice 6D




Now later with inferior Q waves and T wave inversion
The Peaked T in V2 is an analogy of posterior T inversion

Practice 7




Acute anterolateral MI, old enough for Q waves
But T waves still upright and ST segments still up somewhat

Practice 7B



Slightly later, with now some evolutionary terminal T inversion
Notice ST elevation looks worse, but that is due to double-standard


Practice 8



Sinus rhythm with inferior MI and RBBB


An unusual combination, usually pre-existent RBBB

Practice 9




Another acute anterolateral MI, notice ST elevation in I and aVL

Practice 9B




Slightly later, notice the evolutionary T inversion especially in V4 and V5

Practice 10




Acute anterolateral MI and RBBB and leftward axis
Also sinus rhythm with frequent supraventricular ectopy
T inversion in V1 is from the RBBB and not the MI

Practice 10B




Later, new IVCD – QRS looks completely different
Is there 2:1 AV block? Anterior MI and RBBB is risk for AV block!
Complete heart block with ventricular escape?

Practice 11




Sometimes sick patients make for hard tracings
Inferior ST elevation and V1 ST elevation too

Practice 11B




In inferior MI, when ST elevation in V1 is greater than V2,
There is RV MI and not anterior injury

Practice 12




Acute inferior MI. Notice the chest leads have reverse progression, they are right chest leads. ST elevation in V4R indicates RV myocardial infarction

Practice 12B




Inferior MI one hour later, now left chest leads
Complete heart block with narrow junctional escape

Practice 13




Practice 14




Sinus tachycardia, diffuse ST segment elevation
Pericarditis

Practice 15




Close to anterior hyperacute T wave change
But Q in V2-3 already happening and ST elevation also

Practice 15B




3 years later, there are Qs inferiorly as well as anteriorly
Persistent ST and T abnormality

Practice 16




Practice 17




Wellens T wave

Practice 17B




Same patient with Wellens T wave

Practice 18



Close to hyperacute inferior T waves, but some ST elevation
Also reciprocal change in aVL and some elevation in V6

Practice 19



Practice 20

